



**HAL**  
open science

## Alkaline hydrometallurgical approach for steel slag management

Fernando Pereira, Olivier Faure, Veronique Lavastre, Steve Peuble, Frédéric Paran, Jean-Michel Herri, Didier Graillet

### ► To cite this version:

Fernando Pereira, Olivier Faure, Veronique Lavastre, Steve Peuble, Frédéric Paran, et al.. Alkaline hydrometallurgical approach for steel slag management. 2èmes Rencontres Scientifiques: Réhabilitation et Valorisation des Sites et Sols pollués, RV2S - 2nd Scientific Meeting: Rehabilitation of polluted Sites and Soils, RV2S/RSS 2018, Oct 2018, Saint-Etienne, France. Ecole Nationale d'Ingénieurs de Saint-Etienne, 2018, Proceeding: 2èmes Rencontres Scientifiques: Workshop Réhabilitation et valorisation des sites et sols pollués; Proceeding: 2nd Scientific Meeting: Workshop on Rehabilitation of polluted Sites and Soils. emse-03154750

**HAL Id: emse-03154750**

**<https://hal-emse.ccsd.cnrs.fr/emse-03154750>**

Submitted on 1 Mar 2021

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# ALKALINE HYDROMETALLURGICAL APPROACH FOR STEEL SLAG MANAGEMENT



Fernando PEREIRA (1, 3)  
Olivier FAURE (1, 3)  
Véronique LAVASTRE (2, 5)  
Steve PEUBLE (1, 3)  
Frédéric PARAN (1, 3)  
Jean-Michel HERRI (1, 4)  
Didier GRAILLOT (1, 3)







1 - Mines Saint-Étienne - Centre "Sciences des Processus Industriels et Naturels" (SPIN) - "Département Procédés pour l'Environnement et les Géo-ressources" (PEG) - Saint-Étienne (France), 2 - "Université Jean Monnet" (UJM) - Saint-Étienne (France), 3 - CNRS ("Centre National de la Recherche Scientifique") UMR ("Unité Mixte de Recherche") EVS ("Environnement, Ville et Société") 5600 - Lyon (France), 4 - CNRS UMR LGF ("Laboratoire Georges Friedel") - Saint-Étienne (France), 5 - CNRS UMR LMV ("Laboratoire Magmas et Volcans") - Clermont-Ferrand (France) [Contact : [fernando.pereira@mines-stetienne.fr](mailto:fernando.pereira@mines-stetienne.fr)].

## Context

Steel slags are artificial mineral matrices produced by the steel and iron industry. While they are considered as industrial waste, these slag heaps represent an important potential economic resource. The importance of hydrometallurgy in production and recycling of strategic metals cannot be stressed enough. Nevertheless, their use is increasingly under economic pressure since 2014. Clearly, improving or even optimizing these processes is required. In addition, although steel slags are classified as non-hazardous thermal waste, it is well known that most metals they contain are quite toxic to living organisms and pose serious environmental issues. Thus, reusing steel slags appears the option of choice to ensure their sustainable management.

## Experimental

Incorporated into mineral matrices and sometimes committed in the form of ferrites type, strategic metal-based (Cr, W, etc.) non-ferrous oxides have rarely been the subject of specific treatments. When keeping the mineral matrix intact is desired for its subsequent reuse, processing technologies with acid leaching agents are inapplicable and to be avoided. The alkaline earth metals rich composition of steel slags are attractive materials for potential users.



## Conclusions

The innovation was implementing selective unitary operations set on the chemical nature of (EAF-S) non-ferrous metals. The scientific approach consisted in adapting the alkaline middle (use of soda ash as leaching agent) of selective lixiviation and the separation methods in order to optimize the strategic metals extraction rates while choosing conditions and environments that do not present side effects and allow a minimization of the environmental impacts. Soda ash's weak corrosion power and selectivity showed many benefits. Iron and calcium, present in (EAF-S), were poorly leached at the end of alkaline treatment (mineral matrix preservation). Moreover, adding lime in a soda ash solution precipitated the silicates and carbonates thus returned the leaching solution to its initial state.